

7 References

1. Avery J. Brief Report 2004-11. Risky Drinking in South Australia. Population Research and Outcome Studies Unit. 2004. Online [Accessed 6th October 2004] <http://www.dh.sa.gov.au/pehs/PROS/br-risky-drinking04-11.pdf>
2. Australian Bureau of Statistics (ABS) 1993. Estimated resident population by sex and age, states and territories of Australia, June 1987 to June 1992. Cat. no. 3201.0. Canberra
3. Doll R, Payne P & Waterhouse JAH (1966). Cancer incidence in five continents, Vol I. Geneva: UICC, Berlin, Springer.
4. English DR, Holman CDJ, Milne E, Winter MG, Hulse GK, Codde JP et al. 1995. The quantification of drug caused morbidity and mortality in Australia 1995. Canberra: Commonwealth Department of Human Services and Health.
5. International Agency for Research on Cancer (IARC) 2001. GLOBOCAN 2000: cancer incidence, mortality and prevalence worldwide, version 1.0. IARC CancerBase No. 5. Lyon: IARC Press.
6. Jensen OM, Parkin DM, Machennan R & Muir C (eds) 1991. Cancer registration: principles and methods. Lyons: International Agency for Research on Cancer.
7. National Cancer Control Initiative (NCCI) 2003. The 2002 national non-melanoma skin cancer survey: a report by the NCCI Non-melanoma Skin Cancer Working Group. Editor Staples MP. Melbourne: National Cancer Control Initiative.
8. Ridolfo B & Stevenson C 2001. The quantification of drug-caused mortality and morbidity in Australia, 1998. AIHW cat. no. PHE 29. Canberra: AIHW (Drug Statistics Series no. 7).
9. World Health Organization 1992. International statistical classification of diseases and related health problems, 10th revision. Geneva: World Health Organization.